

Calcific Tendinitis

Restoring comfortable movement after treatment for calcific tendinitis.

Kieran Hirpara 4.0



This protocol covers the rehabilitation after **arthroscopic excision of a calcific deposit** with Dr Kieran Hirpara at Mater Private Hospital Rockhampton – keyhole surgery to remove the calcium deposit from the rotator cuff tendon, sometimes combined with a subacromial decompression to make more room for the tendon. Bring this page or its PDF to your first physiotherapy visit so your rehabilitation stays coordinated. Your rehabilitation is progressed individually by your physiotherapist through the phases below, depending on how your shoulder progresses.

If you have any concerns about your wound after surgery, get in touch with the rooms. It is often helpful to take a photo of the wound and email it for review.

If removing the deposit required a rotator cuff repair – the deposit sometimes leaves a defect in the tendon that needs to be stitched closed – your recovery follows rotator cuff repair rules instead, and the [rotator cuff repair protocol](#) takes priority over this page. Dr Hirpara will tell you after the operation if this applies to you.

What to expect

Both pathways are early-movement pathways. Unless the rotator cuff needed a repair, nothing has been stitched that needs months of protection – the aim of treatment is to remove the source of pain, and the aim of rehabilitation is to settle the post-treatment flare, keep the shoulder moving so it does not stiffen, and then rebuild strength.


- **After arthroscopic excision** a sling is provided for comfort only. It is worn briefly – typically days, rarely beyond two weeks – and should be left off as much as possible. You do not need to sleep in it. Recovery to full, unrestricted activity typically takes around three months.

Do not drive while wearing a sling. Driving typically resumes from about two weeks, once you are out of the sling, comfortable, and able to perform an emergency stop safely.

With either treatment, the shoulder can take time to settle completely. Discomfort often improves in stages rather than all at once, and after surgery it can take several months – occasionally up to nine – for the pre-treatment symptoms to fade fully. Steady improvement, not instant comfort, is the expected pattern.

This is keyhole day surgery through several small incisions. The calcific deposit is located within the rotator cuff tendon and removed, and a subacromial decompression is often performed at the same time to give the tendon more room. The dressings can usually come off after about two days, and the wounds are checked at your first post-operative appointment about a week to ten days after surgery.

Phase I – Early movement (weeks 0–2)




Kieran Hirpara © ⓘ 4.0

Open and close hand

Keep your hand and fingers moving by opening and closing them, or by squeezing a soft ball. Start this straight away to keep the hand and arm working while the shoulder settles.

10 times, three times a day




Kieran Hirpara © ⓘ 4.0

Wrist movement

Keep your wrist moving by bending it forwards, backwards and side to side. Do this with the arm out of the sling.

10 times, three times a day




Kieran Hirpara © ⓘ 4.0

Elbow bends

With the upper arm resting by your side, bend the elbow as far as it comfortably goes, then straighten it fully. Only the elbow moves – keep the shoulder relaxed.

10 times, three times a day



Kieran Hirpara © ⓘ 4.0

Assisted overhead stretch (lying)

Lie on your back and use your good arm to help lift the operated arm up above shoulder height as a gentle stretch, then lower it back down with the good arm still helping. It will be uncomfortable at first – that is expected – and it is the key exercise for stopping the shoulder stiffening.

10 repetitions, three times a day

You will wake from surgery with your arm in a sling, but the sling is for comfort only – try to do without it as much as you can, and most people are out of it within days. You can use the arm without restriction below shoulder height from the start. Lifting the arm above shoulder height is allowed and safe, although it will be uncomfortable at first; several times a day, use your good arm to help lift the operated arm above shoulder height as a gentle stretch, so the shoulder does not stiffen. Avoid carrying anything heavier than about two kilograms with the operated arm in these first weeks, as this will be painful. Start your exercises as soon as possible, aiming for ten repetitions of each, three times a day. Take pain relief before your exercises, and use ice for comfort. Do not drive while you are wearing the sling.

For your physiotherapist:

Goals

- Settle post-operative pain and swelling
- Early restoration of range of motion – the priority is preventing stiffness, to which calcific tendinitis patients are prone
- Normal use of the arm below shoulder height

Management

- Sling for comfort only – wean as quickly as comfort allows, typically within days
- Unrestricted active use of the arm below shoulder height from day one
- Active elevation above shoulder height as tolerated
- Passive and active-assisted elevation above shoulder height several times daily (using the other arm) to prevent stiffness

CQ HAND + UPPER LIMB

Dr Kieran Hirpara – Specialist Orthopaedic Surgeon
 Suite 2, Level 1, Mater Private Hospital Rockhampton, 31 Ward Street, The Range, QLD 4700
 Phone 07 4863 6556 · office@cqupperlimb.com.au · cqupperlimb.com.au

- Home exercise program ten repetitions each, three times daily
- Analgesia before exercises; cryotherapy for pain relief as needed


Precautions

- No carrying or lifting beyond approximately two kilograms with the operated arm
- No driving while in the sling

Criteria to progress

- Wound review satisfactory at the first post-operative appointment
- Out of the sling and using the arm comfortably below shoulder height

Phase II – Regaining your range (weeks 2–8)




Kieran Hirpara © ⓘ 4.0

Supported abduction

Cradle the operated forearm with your other arm and lift both out to the side, letting the good arm do most of the work, then lower with control. Progress the range as comfort allows — stretching to firm discomfort is acceptable; forced, severely painful stretching is not.

As guided by your physiotherapist




Kieran Hirpara © ⓘ 4.0

Assisted external rotation with a stick

Lie on your back with the elbow tucked by your side and bent to a right angle. Hold a stick in both hands and use your good arm to turn the operated hand outwards, rotating the shoulder. Restoring this outward rotation is one of the goals of this phase, so keep working it gently a little further each week.

10 times, three times a day




Kieran Hirpara © ⓘ 4.0

Over-door pulley

Sit under an over-door pulley with a handle in each hand. Pull down with your good arm to raise the operated arm overhead as far as it comfortably goes, then lower slowly. Work into a gentle stretch, not sharp pain.

10 times, three to four times a day



Kieran Hirpara © ⓘ 4.0

Scapular setting

Sitting or standing tall, gently draw your shoulder blades back together and down, away from your ears. Hold for a few seconds, then relax. Continue this alongside your stretches to keep the shoulder blade moving well as your range returns.

As guided by your physiotherapist

You will be reviewed in the rooms at about two to three weeks, where your wound and your passive range of motion are checked. The work of this phase is range: progress your forward elevation stretches and add movements out to the side, with physiotherapy guiding the progression. The typical aims are to lift the arm actively to horizontal by six weeks, with the assisted (passive) range – forward, out to the side and in rotation – back to normal by six weeks. Driving can resume from about two weeks once you are out of the sling, comfortable, and able to perform an emergency stop safely.

For your physiotherapist:

Goals

- Active forward flexion and abduction to horizontal by six weeks
- Passive flexion, abduction and external rotation to normal by six weeks
- Independence with daily activities

Management

- Progress passive and active-assisted forward flexion; introduce and progress abduction
- Progress to active range of motion in all planes as comfort allows
- Continue scapular setting and postural work
- Continue analgesia before sessions, and heat or ice around stretching as preferred

Precautions

- Keep lifting light while range is restored; progression remains symptom-guided
- Stretching to firm discomfort is acceptable; forced, severely painful stretching is not


CQ HAND + UPPER LIMB

Dr Kieran Hirpara – Specialist Orthopaedic Surgeon
 Suite 2, Level 1, Mater Private Hospital Rockhampton, 31 Ward Street, The Range, QLD 4700
 Phone 07 4863 6556 · office@cqupperlimb.com.au · cqupperlimb.com.au

Criteria to progress

- Passive range of motion at or near normal
- Active elevation to horizontal or better, with pain settling

Phase III – Strengthening and return to full activity (weeks 8–16)




Kieran Hirpara © ⓘ 4.0

Isometric external rotation

Stand with your elbow tucked at your side and bent to a right angle, the back of your hand against a wall or doorframe. Gently press outwards into it without letting the arm move, hold a few seconds, then relax.

This builds the rotator cuff with no movement — the starting point for strengthening from about eight weeks.

Hold a few seconds, several times; as guided by your physiotherapist




Kieran Hirpara © ⓘ 4.0

Band external rotation

Stand with your elbow tucked into your side and bent to a right angle, holding an elastic band anchored at waist height. Keeping the elbow at your side, rotate the forearm outwards against the band, then return slowly. Progresses on from the isometric work as your strength builds.

2 to 3 sets of 10 to 15, low resistance




Kieran Hirpara © ⓘ 4.0

Low row

Hold a band anchored in front of you at waist height. Keeping the arm fairly straight, pull it back and down towards your hip, squeezing the shoulder blade down and back, then return slowly. Builds the shoulder-blade muscles that support the cuff.

2 to 3 sets of 10 to 15



Kieran Hirpara © ⓘ 4.0

Side-lying external rotation with weight

Lie on your non-operated side with the top elbow bent to a right angle and tucked against your body. Holding a light weight, rotate the forearm up towards the ceiling, then lower slowly. Part of the rotator cuff strengthening from about eight weeks – keep mobility work going alongside it.

Low load, higher repetitions

You will usually be reviewed again at about eight weeks. With your range restored, rehabilitation turns to strengthening the rotator cuff, typically under your physiotherapist's supervision, and to using the arm freely above shoulder height. The aim is full active forward elevation and abduction by about twelve weeks. Recovery from removal of the deposit alone typically takes about three months, after which there are no restrictions; if the rotator cuff needed a repair, recovery is longer – typically around five months – and follows the rotator cuff repair protocol instead. Do not be concerned if some ache persists beyond this point: after this operation it can take up to nine months for the pre-operative symptoms to settle completely, with the trend steadily in the right direction.

For your physiotherapist:

Goals

- Full active forward flexion and abduction by approximately twelve weeks
- Graduated restoration of rotator cuff and scapular strength and endurance
- Return to full, unrestricted activity by approximately three months

Management

- Progressive rotator cuff strengthening from eight weeks – isometrics progressing to elastic-band and light-weight work, low load and higher repetitions
- Progress active use of the arm above shoulder height
- Advance gym-, work- and sport-specific loading as tolerated through weeks twelve to sixteen

CQ HAND + UPPER LIMB

Dr Kieran Hirpara – Specialist Orthopaedic Surgeon
 Suite 2, Level 1, Mater Private Hospital Rockhampton, 31 Ward Street, The Range, QLD 4700
 Phone 07 4863 6556 · office@cqupperlimb.com.au · cqupperlimb.com.au

Precautions

- Strengthening should not come at the cost of range – continue mobility work throughout
- Build heavy and overhead loading gradually; a flare of pain means easing back a step

Criteria to progress

- Full active range of motion with strength returning and symptoms continuing to settle
- Discharge from routine follow-up when progressing well, typically from around eight to sixteen weeks

After your protocol

The phases above are adapted from published patient guidance and rehabilitation protocols for this procedure – The London Shoulder Partnership’s calcific deposit excision rehabilitation protocol, patient guidance from ShoulderDoc (UK) on surgery for calcific tendinitis, and Dr Kevin Ko’s patient guide to arthroscopic excision. The week ranges are typical rather than fixed, and your ongoing rehabilitation is guided individually by your physiotherapist, working with the practice, based on how your shoulder recovers. This page works alongside the practice’s general recovery advice – see [managing post-operative pain](#) and [wound care](#). For the condition itself and how these treatments work, see [calcific tendinitis](#). The evidence behind this protocol – natural history, barbotage, and the surgical-excision literature – is summarised in the evidence section, available as a PDF from the top of this page.