

Shoulder Arthroplasty for Acute Proximal Humerus Fracture

title: "Shoulder Arthroplasty for Acute Proximal Humerus Fracture" slug: proximal-humerus-arthroplasty region: shoulder audience: patient mesh_terms: ["Shoulder Fractures"] article_count: 733 model_used: qwen3.5-35b-a3b-q8 generated_at: '2026-05-18T14:19:37+00:00' key_articles: - title: "Shoulder hemiarthroplasty for acute fractures of the proximal humerus: A minimum five-year follow-up" ref_num: 1 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2007.06.025 year: 2008 - title: "How the greater tuberosity affects clinical outcomes after reverse shoulder arthroplasty for proximal humeral fractures" ref_num: 2 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2018.05.030 year: 2018 - title: "Technique and clinical results of a new intramedullary support nail and plate system for fixation of 3- or 4- part proximal humeral fractures in older adults" ref_num: 3 evidence_tier: paper evidence_level: 3 doi: 10.1186/s12891-022-05998-z year: 2022 - title: "Mortality after inpatient stay for proximal humeral fractures" ref_num: 4 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2019.05.030 year: 2020 - title: "Minimum 2-year outcomes of reverse total shoulder arthroplasty for fracture: how does acute arthroplasty compare with salvage?" ref_num: 5 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2021.06.020 year: 2022 - title: "Functional outcomes of reverse shoulder arthroplasty compared with hemiarthroplasty for acute proximal humeral fractures" ref_num: 6 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2012.03.006 year: 2013 - title: "Fracture line morphology of complex proximal humeral fractures" ref_num: 7 evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jse.2017.05.014 year: 2017 - title: "Improvement in shoulder rotation in complex shoulder fractures treated by reverse shoulder arthroplasty" ref_num: 8 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2012.03.011 year: 2013 - title: "Ten-year follow-up of stemmed hemiarthroplasty for acute proximal humeral fractures" ref_num: 9 evidence_tier: paper evidence_level: 3 doi: 10.1302/0301-620x.103b6.bjj-2020-1753.r1 year: 2021 - title: "Readmissions, revisions, and mortality after treatment for proximal humeral fractures in three large states" ref_num: 10 evidence_tier: paper evidence_level: 3 doi: 10.1186/s12891-019-2812-9 year: 2019 - title: "Consensus statement on the treatment of proximal humerus fractures: a Delphi approach by the Neer Circle of the American Shoulder and Elbow Surgeons" ref_num: 11 evidence_tier: paper evidence_level: 5 doi: 10.1016/j.jse.2024.12.005 year: 2025 - title: "Complications after non-surgical management of proximal humeral fractures: a systematic review of terms and definitions" ref_num: 12 evidence_tier: paper evidence_level: 1 doi: 10.1186/s12891-019-2459-6 year: 2019 - title: "Functional Outcome After Nonoperative Treatment of a Proximal Humeral Fracture in Adults" ref_num: 13 evidence_tier: paper evidence_level: 1 doi: 10.2106/jbjs.20.02018 year: 2021 - title: "Comparison of Hemiarthroplasty and Reverse Arthroplasty for Treatment of Proximal Humeral Fractures" ref_num: 14 evidence_tier: paper evidence_level: 2 doi: 10.2106/jbjs.l.01115 year: 2013 - title: "Morbidity and mortality of

fragility proximal humerus fractures: a retrospective cohort study of patients presenting to a level one trauma center” ref_num: 15 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2022.03.006 year: 2022 - title: “Fragility of randomized controlled trials on treatment of proximal humeral fracture” ref_num: 16 evidence_tier: paper evidence_level: 2 doi: 10.1016/j.jse.2022.01.141 year: 2022 - title: “Isolated avulsion fracture of the lesser tuberosity of the humerus in adolescents. A report of two cases.” ref_num: 17 evidence_tier: paper evidence_level: 4 doi: 10.2106/00004623-199509000-00020 year: 1995 - title: “Hospital Readmissions After Surgical Treatment of Proximal Humerus Fractures: Is Arthroplasty Safer Than Open Reduction Internal Fixation?” ref_num: 19 evidence_tier: paper evidence_level: 3 doi: 10.1007/s11999-014-3613-y year: 2014 - title: “One Versus 3-Week Immobilization Period for Nonoperatively Treated Proximal Humeral Fractures” ref_num: 20 evidence_tier: paper evidence_level: 2 doi: 10.2106/jbjs.20.02137 year: 2021 - title: “In-hospital Complications Are More Likely to Occur After Reverse Shoulder Arthroplasty Than After Locked Plating for Proximal Humeral Fractures” ref_num: 21 evidence_tier: paper evidence_level: 3 doi: 10.1097/corr.0000000000001776 year: 2021 - title: “A new clinical sign to detect radiologically occult greater tuberosity fractures” ref_num: 22 evidence_tier: paper evidence_level: 2 doi: 10.1016/j.jse.2023.07.044 year: 2024 - title: “Proximal Humeral Fracture Treatment in Adults” ref_num: 23 evidence_tier: paper evidence_level: 5 doi: 10.2106/jbjs.l.01293 year: 2014 - title: “Proximal Humeral Fractures: “Damned If You Operate, and Damned If You Don’t”” ref_num: 24 evidence_tier: paper evidence_level: 5 doi: 10.2106/jbjs.19.01109 year: 2019 - title: “Mortality after a proximal humeral fracture” ref_num: 25 evidence_tier: paper evidence_level: 3 doi: 10.1302/0301-620x.102b11.bjj-2020-0627.r1 year: 2020 - title: “Hemiarthroplasty versus reverse shoulder arthroplasty for treatment of proximal humeral fractures: a meta-analysis” ref_num: 26 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2015.08.030 year: 2016 - title: “Short-term Complications for Proximal Humerus Fracture Surgery Have Decreased: An Analysis of the National Surgical Quality Improvement Program Database” ref_num: 27 evidence_tier: paper evidence_level: 3 doi: 10.1097/corr.0000000000002391 year: 2022 - title: “The role of greater tuberosity healing in reverse shoulder arthroplasty: a finite element analysis” ref_num: 28 evidence_tier: paper evidence_level: 5 doi: 10.1016/j.jse.2019.07.022 year: 2020 - title: “Does objective shoulder impairment explain patient-reported functional outcome? A study of proximal humerus fractures” ref_num: 29 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2010.06.005 year: 2011 - title: “Neer Award 2006: Biomechanical assessment of inferior tuberosity placement during hemiarthroplasty for four-part proximal humeral fractures” ref_num: 30 evidence_tier: paper evidence_level: 5 doi: 10.1016/j.jse.2007.06.017 year: 2008 - title: “Evaluation of the Constant score: which is the method to assess the objective strength?” ref_num: 35 evidence_tier: paper evidence_level: 3 doi: 10.1186/s12891-019-2795-6 year: 2019 - title: “Does fracture of the dominant shoulder have any effect on functional and quality of life outcome compared with the nondominant shoulder?” ref_num: 37 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2014.10.006 year: 2015 - title: “The assessment of scapular radiographs” ref_num: 38 evidence_tier: paper evidence_level: 4 doi: 10.1302/0301-620x.95b8.30631 year: 2013 - title: “Altered Glenohumeral Biomechanics in Proximal Humeral Fracture Malunion” ref_num: 40 evidence_tier: paper evidence_level: 5 doi: 10.5435/jaaos-d-20-00555 year: 2020 - title: “Effects of shoulder position on axillary nerve positions during the split lateral deltoid approach” ref_num: 43 evidence_tier: paper evidence_level: 5 doi: 10.1016/j.jse.2008.12.001 year: 2009 - title: “A morphovolumetric study of head malposition in proximal humeral fractures based on 3-dimensional computed tomography scans: the control volume theory” ref_num: 44 evidence_tier: paper evidence_level: 5 doi: 10.1016/j.jse.2017.12.004 year: 2018 - title: “Cost-Minimization Analysis and Treatment Trends of Surgical and Nonsurgical Treatment of Proximal Humerus Fractures” ref_num: 46

evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jhsa.2020.03.022 year: 2020 - title: "Reverse shoulder arthroplasty or nothing for patients with displaced proximal humeral fractures: a randomized controlled trial" ref_num: 47 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2024.02.023 year: 2024 - title: "Operative versus non-operative treatment for 2-part proximal humerus fracture: A multicenter randomized controlled trial" ref_num: 48 evidence_tier: paper evidence_level: 1 doi: 10.1371/journal.pmed.1002855 year: 2019 - title: "Classification of proximal humeral fractures based on a pathomorphologic analysis" ref_num: 49 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2015.08.006 year: 2016 - title: "Evaluation and Management of Pediatric Proximal Humerus Fractures" ref_num: 50 evidence_tier: paper evidence_level: 5 doi: 10.5435/jaaos-d-14-00033 year: 2015 - title: "Contemporary Management of Proximal Humeral Fractures" ref_num: 51 evidence_tier: paper evidence_level: 5 doi: 10.5435/jaaos-d-24-01073 year: 2025 - title: "Locking plate versus nonsurgical treatment for proximal humeral fractures: better midterm outcome with nonsurgical treatment" ref_num: 52 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2011.01.025 year: 2011 - title: "Osteoporosis does not affect bone mineral density change in the proximal humerus or the functional outcome after open reduction and internal fixation of unilateral displaced 3- or 4-part fractures at 12-month follow-up" ref_num: 53 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2022.07.008 year: 2023 - title: "Reverse Shoulder Arthroplasty to Treat Proximal Humerus Fracture Sequelae: A Review" ref_num: 54 evidence_tier: paper evidence_level: 5 doi: 10.5435/jaaos-d-23-00740 year: 2024 - title: "Path analysis of factors for functional outcome at one year in 463 proximal humeral fractures" ref_num: 55 evidence_tier: paper evidence_level: 2 doi: 10.1016/j.jse.2011.06.009 year: 2011 - title: "A comprehensive classification of proximal humeral fractures: HGLS system" ref_num: 56 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2012.09.018 year: 2013 - title: "CORR Insights®: What Factors Are Associated With Poor Shoulder Function and Serious Complications After Internal Fixation of Three-part and Four-part Proximal Humerus Fracture-dislocations?" ref_num: 57 evidence_tier: paper evidence_level: 5 doi: 10.1097/corr.0000000000002242 year: 2022 - title: "Four-segment classification of proximal humeral fractures revisited: A multicenter study on 509 cases" ref_num: 58 evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jse.2009.01.018 year: 2009 - title: "Outcome measures in the management of proximal humeral fractures: a systematic review of their use and psychometric properties" ref_num: 59 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2010.10.028 year: 2011 - title: "Evaluation of the Neer System of Classification of Proximal Humeral Fractures with Computerized Tomographic Scans and Plain Radiographs*" ref_num: 60 evidence_tier: paper evidence_level: 4 doi: 10.2106/00004623-199609000-00012 year: 1996 - title: "Outcomes after percutaneous reduction and fixation of proximal humeral fractures" ref_num: 62 evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jse.2006.09.006 year: 2007 - title: "Long-term outcomes of reverse shoulder arthroplasty versus nonoperative treatment for 3- or 4-part proximal humerus fractures in elderly patients: results from a prior randomized clinical trial" ref_num: 63 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2024.09.032 year: 2025 - title: "Is there still a place for anatomic hemiarthroplasty in patients with high functional demands in primary, nonreconstructable proximal humeral fractures? A clinical and radiographic assessment" ref_num: 65 evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jse.2023.02.118 year: 2023 - title: "Clinical outcomes of reverse total shoulder arthroplasty for elective indications versus acute 3- and 4-part proximal humeral fractures: a systematic review and meta-analysis" ref_num: 66 evidence_tier: paper evidence_level: 1 doi: 10.1016/j.jse.2021.07.014 year: 2022 - title: "Hemiarthroplasty for Humeral Four-part Fractures for Patients 65 Years and Older: A Randomized Controlled Trial" ref_num: 67 evidence_tier: paper evidence_level: 1 doi: 10.1007/s11999-012-2531-0 year: 2012 - title: "The effect of medial calcar support on proximal humeral fractures treated with locking plates" ref_num: 71 evidence_tier: paper evidence_level: 3 doi:

10.1186/s13018-022-03337-5 year: 2022 - title: "Missed fractures of the greater tuberosity" ref_num: 72 evidence_tier: paper evidence_level: 4 doi: 10.1186/s12891-018-2225-1 year: 2018 - title: "Coronal displacement in proximal humeral fractures: correlation between shoulder radiographic and computed tomography scan measurements" ref_num: 73 evidence_tier: paper evidence_level: 3 doi: 10.1016/j.jse.2016.05.016 year: 2017 - title: "CORR Insights®: 3D-printed Handheld Models Do Not Improve Recognition of Specific Characteristics and Patterns of Three-part and Four-part Proximal Humerus Fractures" ref_num: 74 evidence_tier: paper evidence_level: 5 doi: 10.1097/corr.0000000000002017 year: 2021 - title: "Detection, classification, and characterization of proximal humerus fractures on plain radiographs" ref_num: 76 evidence_tier: paper evidence_level: 3 doi: 10.1302/0301-620x.106b11.bjj-2024-0264.r1 year: 2024 - title: "Missed posterior dislocation of the shoulder after intramedullary fixation of humeral fractures: A report of three cases" ref_num: 79 evidence_tier: paper evidence_level: 4 doi: 10.1016/j.jse.2008.10.020 year: 2009 - title: "Does time to surgery affect patient-reported outcome in proximal humeral fractures? A subanalysis of the PROFHER randomized clinical trial" ref_num: 80 evidence_tier: paper evidence_level: 1 doi: 10.1302/0301-620x.102b1.bjj-2020-0546.r1 year: 2020 synthesis_version: "v2" verifier_status: skipped

Overview

- Patients undergoing arthroplasty for acute proximal humeral fractures may achieve satisfactory long-term pain relief, though overall shoulder motion results are less predictable [1].
- In elderly patients undergoing reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes [2].
- Clinical results at 1-year follow-up confirm the advantage of applying a new intramedullary support nail and plate system to 3- or 4-part proximal humeral fractures in older patients [3].
- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- Patients who undergo initial nonoperative management have worse functional outcomes and higher complication rates than those who undergo acute reverse total shoulder arthroplasty (rTSA) for proximal humeral fractures [5].
- Patients with acute proximal humeral fractures who undergo reverse shoulder arthroplasty appear to achieve superior 5-year functional outcomes compared with patients who undergo hemiarthroplasty [6].
- The study cited represents the largest long-term follow-up of acute proximal humeral fractures treated with hemiarthroplasty [9].
- In most studies of proximal humeral fractures, only 1 or 2 patients experiencing an alternative outcome or lost to follow-up would change the conclusions for the dichotomous outcome studied [16].
- The increased in-hospital risk for major adverse events and surgical complications may moderate enthusiasm for reverse total shoulder arthroplasty (RTSA) for proximal humeral fractures in patients 65 years and older [21].

- Available literature suggests that reverse shoulder arthroplasty performed to address complex proximal humeral fractures might result in more favorable clinical outcomes than hemiarthroplasty performed for the same indication [26].
- Reverse total shoulder arthroplasty (RTSA) performed for acute 3- and 4-part proximal humeral fractures yields overall worse clinical outcomes and active range of motion compared with RTSA performed for elective indications [66].
- No clear benefits were observed in treating patients 65 years or older with four-part fractures of the proximal humerus with either hemiarthroplasty or nonoperative treatment [67].

Anatomy & Pathophysiology

- Shoulder rotational ability is improved by systematically repairing the tuberosities around the implant, provided their consolidation is anatomic [8].
- Greater tuberosity healing does not seem to impact reverse shoulder arthroplasty biomechanics during abduction or forward flexion [28].
- Greater tuberosity healing affects reverse shoulder arthroplasty biomechanics during external rotation [28].
- With minimal and moderate amounts of glenohumeral abduction, glenohumeral joint forces are significantly displaced superiorly [30].
- Varus and antecurvatum proximal humerus deformities as small as 15 degrees were associated with statistically significant alterations in glenohumeral joint mechanics [40].
- The control volume is an important anatomic and functional area of the proximal humerus [44].
- Vertical abduction has the greatest effect on axillary nerve position during the split lateral deltoid approach [43].
- Horizontal glenohumeral forward flexion and humeral rotation have little effect on axillary nerve position during the split lateral deltoid approach [43].
- The study demonstrates variability in the glenopolar angle with increased AP rotational offset of the shoulder radiograph [38].
- The study reveals inaccuracies in glenopolar angle measurement even at an institution with an established protocol [38].
- Range of motion and strength thresholds can identify subjects with normal shoulder function [29].
- The authors recommend performing the measurement of objective strength at the insertion of the deltoid muscle in a 90° abduction position in the scapula plane [35].
- Dominance of the affected shoulder has no influence on functional and quality of life outcome compared with the nondominant shoulder [37].
- Dominance of the affected shoulder should not be used to make treatment decisions [37].

Classification

- The Neer classification system covers 98% of all proximal humeral fractures and is appropriate for clinical practice [58].
- Classifications of proximal humeral fractures using the Neer system based on CT scans and plain radiographs are not very reliable or reproducible due to difficulty in determining which segments are fractured [60].
- The HGLS classification is a reliable method of describing fractures of the proximal humerus compared with the Neer and AO systems [56].
- A new classification system with emphasis on the qualitative aspects of proximal humeral fractures showed high reliability when based on a standardized imaging protocol including computed tomography scans [49].
- Consensus when managing proximal humerus fractures is limited to specific scenarios, whereas lack of consensus still exists in others [11].

Clinical Presentation

- Patients undergoing arthroplasty for acute proximal humerus fractures may achieve satisfactory long-term pain relief, though overall shoulder motion results are less predictable [1].
- In elderly patients undergoing reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes [2].
- Clinical results at 1-year follow-up confirm the advantage of applying a new intramedullary support nail and plate system to 3- or 4-part proximal humeral fractures in older patients [3].
- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- Patients undergoing initial nonoperative management have worse functional outcomes and higher complication rates than those undergoing acute reverse total shoulder arthroplasty for proximal humeral fractures [5].
- Fractures of the proximal humerus follow characteristic patterns [7].
- A majority of patients with proximal humeral fractures undergo non-operative treatment [10].
- Consensus on managing proximal humerus fractures is limited to specific scenarios, while a lack of consensus exists in others [11].
- There is significant heterogeneity in the terminology and definitions used to describe complications following non-surgical management of proximal humeral fractures [12].
- Nonoperative treatment of proximal humeral fractures produces considerable variation in shoulder-specific and general health outcomes at 1 year, with a substantial proportion of patients having poor perceived functional outcomes [13].
- Mortality at 1 year for fragility proximal humerus fractures is universally high regardless of risk factors [15].

- The majority of unplanned hospital readmissions after surgical treatment of proximal humerus fractures are associated with medical diagnoses [19].
- In patients presenting with a traumatic shoulder injury and normal radiographs, the anterior bruise sign (ABS) is a highly sensitive and specific clinical aid to identify occult greater tuberosity fractures [22].
- Most proximal humeral fractures in elderly patients can be treated nonoperatively with good functional outcomes [23].
- Patients sustaining a proximal humeral fracture have a significantly higher risk of mortality up to one year after the injury compared with the general population [25].
- Most pediatric patients with proximal humerus fractures have favorable results, and complications are infrequent [50].
- Reverse shoulder arthroplasty is a powerful tool for managing proximal humerus fracture sequelae when joint-preserving options are not optimal, provided there is careful management of the tuberosities and understanding of associated pearls and pitfalls [54].
- Prevention of local complications, particularly those leading to severe varus deviation, appears essential to improve shoulder function after a proximal humeral fracture [55].
- Factors associated with poor results after internal fixation of three-part and four-part proximal humerus fracture-dislocations include being a woman, four-part fracture dislocation, and absence of metaphyseal head extension [57].
- A wide range of outcome measures are used in proximal humeral fracture studies, but there is limited evidence regarding their psychometric properties in this specific population [59].

Investigations

- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- A majority of patients with proximal humeral fractures undergo non-operative treatment [10].
- Fractures of the proximal humerus follow characteristic patterns [7].
- Despite a delayed diagnosis of more than one year, osteotomy and realignment of a displaced lesser tuberosity fracture can be successful and enhance overall shoulder function [17].
- Undisplaced greater tuberosity fractures can be managed non-operatively with good results [72].
- Patients with persistent post-traumatic shoulder pain and limitation of function warrant MRI investigation to identify occult greater tuberosity fractures [72].
- In patients presenting with a traumatic shoulder injury with normal radiographs, the anterior bruise sign (ABS) is a highly sensitive and specific clinical aid to identify patients with an occult greater tuberosity fracture [22].
- There is relevant variability in displacement measurements between shoulder radiographs and CT scans in the coronal plane [73].

- Nearly 30% of cases suggesting surgical treatment on radiographs are reclassified for conservative treatment based on CT findings [73].
- The inherent nature of medial comminution of proximal humeral fracture may lead to inferior radiographic outcomes [71].
- Routine use of 3D-printed models may not be beneficial for classifying proximal humeral fracture patterns beyond the information gained from currently available imaging modalities [74].
- The routine use of 3D-printed models should be avoided as the sole determinant for recommending surgical intervention at this time [74].
- Convolutional neural networks (CNNs) proficiently rule out proximal humerus fractures on plain radiographs [76].
- Missed posterior dislocation of the shoulder after intramedullary fixation of proximal humeral fractures is an extremely rare injury that can be missed due to inadequate initial and postoperative x-ray images and incorrect interpretation [79].
- Consensus when managing proximal humerus fractures is limited to specific scenarios, whereas lack of consensus still exists in others [11].

Treatment

- Patients undergoing shoulder hemiarthroplasty for acute proximal humerus fractures may achieve satisfactory long-term pain relief, though overall shoulder motion results are less predictable [1].
- In elderly patients undergoing reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes [2].
- Clinical results at 1-year follow-up confirm the advantage of applying a new intramedullary support nail and plate system to 3- or 4-part proximal humeral fractures in older patients [3].
- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- Patients undergoing initial nonoperative management have worse functional outcomes and higher complication rates than those undergoing acute reverse total shoulder arthroplasty (rTSA) for proximal humeral fractures [5].
- Patients with acute proximal humeral fractures who undergo reverse shoulder arthroplasty appear to achieve superior 5-year functional outcomes compared with patients who undergo hemiarthroplasty [6].
- Shoulder rotational ability is improved by systematically repairing the tuberosities around the implant in complex shoulder fractures treated by reverse shoulder arthroplasty, provided their consolidation is anatomic [8].
- A majority of patients with proximal humeral fractures underwent non-operative treatment [10].
- Significant heterogeneity exists in the terminology and definitions used to describe complications following non-surgical management of proximal humeral fractures [12].

- Nonoperative treatment of proximal humeral fractures produces considerable variation in shoulder-specific and general health outcomes at 1 year, with a substantial proportion of patients having poor perceived functional outcomes [13].
- Primary shoulder hemiarthroplasty for proximal humeral fracture is associated with satisfactory prosthetic survival at an average of 6.3 years [14].
- Short and long periods of immobilization yield similar results for nonoperatively treated proximal humeral fractures, independent of the fracture pattern [20].
- Most proximal humeral fractures in elderly patients can be treated nonoperatively with good functional outcomes [23].
- Available literature suggests that reverse shoulder arthroplasty performed to address complex proximal humeral fractures might result in more favorable clinical outcomes than hemiarthroplasty performed for the same indication [26].
- Nonsurgical management of proximal humerus fractures decreased during the study period [46].
- Treatment with reverse shoulder arthroplasty provides superior functional outcomes compared with conservative treatment for patients presenting with an acute proximal humeral fracture [47].
- There is no significant difference in clinical outcomes at 2 years between surgery and non-operative treatment in patients 60 years of age or older with displaced 2-part fractures of the proximal humerus [48].
- Nonsurgical management of proximal humerus fractures demonstrates successful outcomes and union rates greater than 90% [51].
- Nonsurgical treatment should have a more prominent role in the treatment of proximal humeral fractures compared to locking plate fixation [52].
- Osteoporosis may not be regarded as a contraindication for open reduction and internal fixation of unilateral displaced 3- or 4-part fractures, as shoulder function was restored to preinjury levels for most patients at 12-month follow-up [53].
- Percutaneous treatment of selected proximal humeral fractures results in predictable union and good clinical results with a low rate of complications [62].
- With narrow indications, use of a specific fracture stem and adequate tuberosity management, successful radiographic and functional results are presented after a mean follow-up of 4.8 years after hemiarthroplasty for primary nonreconstructable humeral head fractures [65].

Complications

- Patients undergoing arthroplasty for acute proximal humeral fractures may achieve satisfactory long-term pain relief, but overall shoulder motion results are less predictable [1].
- In elderly patients undergoing reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes [2].
- Clinical results at 1-year follow-up confirm the advantage of applying a new intramedullary support nail and plate system to 3- or 4-part proximal humeral fractures in older patients [3].

- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- Patients undergoing initial nonoperative management have worse functional outcomes and higher complication rates than those undergoing acute reverse total shoulder arthroplasty (rTSA) for proximal humeral fractures [5].
- Patients with acute proximal humeral fractures who undergo reverse shoulder arthroplasty (RSA) appear to achieve superior 5-year functional outcomes compared with patients who undergo hemiarthroplasty [6].
- Fractures of the proximal humerus follow characteristic patterns [7].
- Primary shoulder hemiarthroplasty for proximal humeral fracture is associated with satisfactory prosthetic survival at an average of 6.3 years [14].
- Mortality at 1 year for fragility proximal humerus fractures is universally high regardless of risk factors [15].
- The majority of unplanned hospital readmissions after surgical treatment of proximal humerus fractures are associated with medical diagnoses [19].
- In-hospital complications are more likely to occur after reverse shoulder arthroplasty than after locked plating for proximal humeral fractures [21].
- Surgery for complex proximal humeral fractures leads to overall good long-term outcomes but is associated with high overall complication and reoperation rates [24].
- Patients sustaining a proximal humeral fracture have a significantly higher risk of mortality up to one year after the injury compared with the general population [25].
- Short-term complication rates for fixation and arthroplasty alike have decreased compared with recent historic norms [27].

Recovery

- Patients undergoing arthroplasty for acute proximal humerus fractures may achieve satisfactory long-term pain relief, though overall shoulder motion results are less predictable [1].
- In elderly patients undergoing reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes [2].
- Clinical results at 1-year follow-up confirm the advantage of applying a new intramedullary support nail and plate system to 3- or 4-part proximal humeral fractures in older patients [3].
- Elderly patients requiring admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age [4].
- Patients who undergo initial nonoperative management have worse functional outcomes and higher complication rates than those who undergo acute reverse total shoulder arthroplasty (rTSA) for proximal humeral fractures [5].
- Patients with acute proximal humeral fractures who undergo reverse shoulder arthroplasty appear to achieve superior 5-year functional outcomes compared with patients who undergo hemiarthroplasty [6].

- Fractures of the proximal humerus follow characteristic patterns [7].
- This study represents the largest long-term follow-up of acute proximal humeral fractures treated with hemiarthroplasty [9].
- Nonoperative treatment of proximal humeral fractures produces considerable variation in shoulder-specific and general health outcomes at 1 year, with a substantial proportion of patients having poor perceived functional outcomes [13].
- Primary shoulder hemiarthroplasty for proximal humeral fracture is associated with satisfactory prosthetic survival at an average of 6.3 years [14].
- Mortality at 1 year for fragility proximal humerus fractures is universally high regardless of risk factors [15].
- In most studies of proximal humeral fractures, only 1 or 2 patients experiencing an alternative outcome or lost to follow-up would change the conclusions for the dichotomous outcome studied [16].
- Despite a delayed diagnosis of more than one year, osteotomy and realignment of a displaced lesser tuberosity fracture was successful and enhanced overall shoulder function in two adolescent patients [17].
- Short and long periods of immobilization yield similar results for nonoperatively treated proximal humeral fractures, independent of the fracture pattern [20].
- Surgery for complex proximal humeral fractures leads to overall good long-term outcomes despite high overall complication and reoperation rates [24].
- The increasing utilization of reverse total shoulder arthroplasty (RTSA) and decreasing short-term complication rates for fixation and arthroplasty represent a substantial change compared with recent historic norms in the management of proximal humerus fractures [27].
- Long-term treatment with reverse shoulder arthroplasty (RSA) for displaced 3- or 4-part proximal humerus fractures provides better functional outcomes compared to nonoperative treatment, a difference attributed to the deterioration of functional outcomes of the nonoperative treatment over time [63].
- Timing of surgery did not affect Oxford Shoulder Score at any stage of follow-up, irrespective of age or fracture type [80].

Key Evidence

- [L3] Patients undergoing arthroplasty as treatment of an acute fracture of the proximal humerus may achieve satisfactory long-term pain relief; however, the result for overall shoulder motion is less predictable. ([10.1016/j.jse.2007.06.025](https://doi.org/10.1016/j.jse.2007.06.025))
- [L3] In elderly patients who have undergone a reverse shoulder arthroplasty for acute proximal humeral fractures, anatomic tuberosity healing improves objective and subjective outcomes. ([10.1016/j.jse.2018.05.030](https://doi.org/10.1016/j.jse.2018.05.030))
- [L3] Clinical results at 1-year follow-up confirmed the advantage of applying it to 3- or 4-part proximal humeral fractures in older patients. ([10.1186/s12891-022-05998-z](https://doi.org/10.1186/s12891-022-05998-z))

- [L3] Elderly patients who require admission after sustaining a proximal humeral fracture are frail and subject to a greater-than-average risk of mortality for their age. ([10.1016/j.jse.2019.05.030](#))
- [L3] Patients who undergo initial periods of nonoperative management have worse functional outcomes and higher complication rates than those who undergo acute rTSA for proximal humeral fractures. ([10.1016/j.jse.2021.06.020](#))
- [L3] Patients with acute proximal humeral fractures who undergo RSA appear to achieve superior 5-year functional outcomes compared with patients who undergo hemiarthroplasty. ([10.1016/j.jse.2012.03.006](#))
- [L4] Fractures of the proximal humerus follow characteristic patterns. ([10.1016/j.jse.2017.05.014](#))
- [L3] Shoulder rotational ability is improved by systematically repairing the tuberosities around the implant, provided their consolidation is anatomic. ([10.1016/j.jse.2012.03.011](#))
- [L3] This is the largest long-term follow-up study of acute proximal humeral fractures treated with hemiarthroplasty. ([10.1302/0301-620x.103b6.bjj-2020-1753.r1](#))
- [L3] A majority of patients with proximal humeral fractures underwent non-operative treatment. ([10.1186/s12891-019-2812-9](#))
- [L5] Consensus when managing proximal humerus fractures is limited to specific scenarios, whereas lack of consensus still exists in others. ([10.1016/j.jse.2024.12.005](#))
- [L1] This systematic review highlights significant heterogeneity in the terminology and definitions used to describe complications following non-surgical management of proximal humeral fractures, calling for standardized definitions to improve evidence synthesis. ([10.1186/s12891-019-2459-6](#))
- [L1] Nonoperative treatment of proximal humeral fractures produces considerable variation in shoulder-specific and general health outcomes at 1 year, and a substantial proportion of patients have poor perceived functional outcomes. ([10.2106/jbjs.20.02018](#))
- [L2] Primary shoulder hemiarthroplasty for proximal humeral fracture is associated with satisfactory prosthetic survival at an average of 6.3 years. ([10.2106/jbjs.l01115](#))
- [L3] Mortality at 1 year for fragility proximal humerus fractures is universally high regardless of risk factors. ([10.1016/j.jse.2022.03.006](#))
- [L2] In most studies of proximal humeral fractures, only 1 or 2 patients experiencing an alternative outcome or lost to follow-up would change the conclusions for the dichotomous outcome studied. ([10.1016/j.jse.2022.01.141](#))
- [L4] Despite a delayed diagnosis of more than one year, osteotomy and realignment of the displaced fracture of the lesser tuberosity was successful and enhanced the overall function of the shoulder in these two patients. ([10.2106/00004623-199509000-00020](#))
- [L3] As the majority of unplanned hospital readmissions were associated with medical diagnoses, it is important to consider patient medical comorbidities before surgical treatment of proximal humerus fractures and during the postoperative care phase. ([10.1007/s11999-014-3613-y](#))
- [L2] Short and long periods of immobilization yield similar results for nonoperatively treated proximal humeral fractures, independent of the fracture pattern. ([10.2106/jbjs.20.02137](#))

- [L3] The increased in-hospital risk for major adverse events and surgical complications may moderate the enthusiasm associated with RTSA for proximal humeral fractures in patients 65 years and older. ([10.1097/corr.0000000000001776](#))
- [L2] In patients presenting with a traumatic shoulder injury with normal radiographs, the anterior bruise sign (ABS) is a highly sensitive and specific clinical aid to identify patients with an occult greater tuberosity fracture. ([10.1016/j.jse.2023.07.044](#))
- [L5] Most proximal humeral fractures in elderly patients can be treated nonoperatively with good functional outcomes. ([10.2106/jbjs.1.01293](#))
- [L5] Surgery for complex proximal humeral fractures leads to overall good long-term outcomes with high overall complication and reoperation rates. ([10.2106/jbjs.19.01109](#))
- [L3] Compared with the general population, patients sustaining a proximal humeral fracture have a significantly higher risk of mortality up to one year after the injury. ([10.1302/0301-620x.102b11.bjj-2020-0627.r1](#))
- [L1] The available literature suggests that reverse shoulder arthroplasty performed to address complex proximal humeral fractures might result in more favorable clinical outcomes than hemiarthroplasty performed for the same indication. ([10.1016/j.jse.2015.08.030](#))
- [L3] The increasing utilization of RTSA and decreasing short-term complication rates for fixation and arthroplasty alike represent a substantial change compared even with recent historic norms in the management of proximal humerus fractures. ([10.1097/corr.0000000000002391](#))
- [L5] Greater tuberosity healing does not seem to impact reverse shoulder arthroplasty biomechanics during abduction or forward flexion; however, it does affect biomechanics during external rotation. ([10.1016/j.jse.2019.07.022](#))
- [L3] Range of motion and strength thresholds can identify subjects with normal shoulder function. ([10.1016/j.jse.2010.06.005](#))
- [L5] With minimal and moderate amounts of glenohumeral abduction, glenohumeral joint forces are significantly displaced superiorly. ([10.1016/j.jse.2007.06.017](#))
- [L3] The authors recommend performing the measurement at the insertion of the deltoid muscle in a 90° abduction position in the scapula plane. ([10.1186/s12891-019-2795-6](#))
- [L3] Dominance of the affected shoulder has no influence and should not be used to make treatment decisions. ([10.1016/j.jse.2014.10.006](#))
- [L4] The study demonstrates variability in the glenopolar angle with increased AP rotational offset of the shoulder radiograph, revealing inaccuracies even at an institution with an established protocol. ([10.1302/0301-620x.95b8.30631](#))
- [L5] Varus and antecurvatum proximal humerus deformities as small as 15 degrees were associated with statistically significant alterations in glenohumeral joint mechanics. ([10.5435/jaaos-d-20-00555](#))
- [L5] Vertical abduction has the greatest effect on axillary nerve position, while horizontal glenohumeral forward flexion and humeral rotation have little effect. ([10.1016/j.jse.2008.12.001](#))
- [L5] The control volume is an important anatomic and functional area of the proximal humerus. ([10.1016/j.jse.2017.12.004](#))

- [L4] Nonsurgical management of proximal humerus fractures decreased during the study period. ([10.1016/j.jhsa.2020.03.022](#))
- [L1] Treatment with reverse shoulder arthroplasty provides superior functional outcomes compared with conservative treatment for patients presenting with an acute proximal humeral fracture. ([10.1016/j.jse.2024.02.023](#))
- [L1] This trial found no significant difference in clinical outcomes at 2 years between surgery and non-operative treatment in patients 60 years of age or older with displaced 2-part fractures of the proximal humerus. ([10.1371/journal.pmed.1002855](#))
- [L3] The new classification system with emphasis on the qualitative aspects of proximal humeral fractures showed high reliability when based on a standardized imaging protocol including computed tomography scans. ([10.1016/j.jse.2015.08.006](#))
- [L5] Most pediatric patients with proximal humerus fractures have favorable results, and complications are infrequent. ([10.5435/jaaos-d-14-00033](#))
- [L5] Treatment for proximal humerus fractures remains controversial, with nonsurgical management demonstrating successful outcomes and union rates greater than 90%. ([10.5435/jaaos-d-24-01073](#))
- [L3] Nonsurgical treatment should have a more prominent role in the treatment of proximal humeral fractures. ([10.1016/j.jse.2011.01.025](#))
- [L1] Shoulder function was restored to preinjury levels for most patients, and osteoporosis may not be regarded as a contraindication for this treatment. ([10.1016/j.jse.2022.07.008](#))
- [L5] Reverse shoulder arthroplasty is a powerful tool for managing proximal humerus fracture sequelae when joint-preserving options are not optimal, provided there is careful management of the tuberosities and understanding of associated pearls and pitfalls. ([10.5435/jaaos-d-23-00740](#))
- [L2] Prevention of local complications, in particular those leading to severe varus deviation, appears essential to improve shoulder function after a proximal humeral fracture. ([10.1016/j.jse.2011.06.009](#))
- [L3] The HGLS classification is a reliable method of describing fractures of the proximal humerus compared with the Neer and AO systems. ([10.1016/j.jse.2012.09.018](#))
- [L5] Surgical treatment of proximal humerus fractures remains far from straightforward, with unpredictable outcomes where factors associated with poor results include being a woman, four-part fracture dislocation, and absence of metaphyseal head extension. ([10.1097/corr.0000000000002242](#))
- [L4] The revised Neer classification covers 98% of all proximal humeral fractures and is appropriate for clinical practice. ([10.1016/j.jse.2009.01.018](#))
- [L1] The review identified a wide range of outcome measures used in proximal humeral fracture studies, but found limited evidence regarding their psychometric properties in this specific population. ([10.1016/j.jse.2010.10.028](#))
- [L4] Classifications of proximal humeral fractures using the Neer system based on CT scans and plain radiographs are not very reliable or reproducible due to difficulty in determining which segments are fractured. ([10.2106/00004623-199609000-00012](#))
- [L4] Percutaneous treatment of selected proximal humeral fractures results in predictable union and good clinical results with a low rate of complications. ([10.1016/j.jse.2006.09.006](#))

- [L1] Long-term treatment with RSA for displaced 3- or 4-part proximal humerus fractures provides better functional outcomes compared to nonoperative treatment, a difference attributed to the deterioration of functional outcomes of the nonoperative treatment over time. ([10.1016/j.jse.2024.09.032](https://doi.org/10.1016/j.jse.2024.09.032))
- [L4] With narrow indications, use of a specific fracture stem and adequate tuberosity management, successful radiographic and functional results are presented after a mean follow-up of 4.8 years after hemiarthroplasty for primary nonreconstructable humeral head fractures. ([10.1016/j.jse.2023.02.118](https://doi.org/10.1016/j.jse.2023.02.118))
- [L1] RTSA performed for acute 3- and 4-part proximal humeral fractures yields overall worse clinical outcomes and active ROM compared with RTSA performed for elective indications. ([10.1016/j.jse.2021.07.014](https://doi.org/10.1016/j.jse.2021.07.014))
- [L1] We observed no clear benefits in treating patients 65 years or older with four-part fractures of the proximal humerus with either hemiarthroplasty or nonoperative treatment. ([10.1007/s11999-012-2531-0](https://doi.org/10.1007/s11999-012-2531-0))
- [L3] This implies that the inherent nature of medial comminution of proximal humeral fracture may lead to inferior radiographic outcomes. ([10.1186/s13018-022-03337-5](https://doi.org/10.1186/s13018-022-03337-5))
- [L4] Undisplaced greater tuberosity fractures can be managed non-operatively with good results, but patients with persistent post-traumatic shoulder pain and limitation of function warrant MRI investigation to identify occult fractures. ([10.1186/s12891-018-2225-1](https://doi.org/10.1186/s12891-018-2225-1))
- [L3] There is relevant variability in displacement measurements between shoulder radiographs and CT scans in the coronal plane, with nearly 30% of cases suggesting surgical treatment on radiographs being reclassified for conservative treatment based on CT findings. ([10.1016/j.jse.2016.05.016](https://doi.org/10.1016/j.jse.2016.05.016))
- [L5] The routine use of 3D-printed models may not be beneficial for classifying proximal humeral fracture patterns beyond the information gained from currently available imaging modalities, and their use as the sole determinant for recommending surgical intervention should be avoided at this time. ([10.1097/corr.0000000000002017](https://doi.org/10.1097/corr.0000000000002017))
- [L3] CNNs proficiently rule out proximal humerus fractures on plain radiographs. ([10.1302/0301-620x.106b11.bjj-2024-0264.r1](https://doi.org/10.1302/0301-620x.106b11.bjj-2024-0264.r1))
- [L4] Missed posterior dislocation of the shoulder after intramedullary fixation of proximal humeral fractures is an extremely rare injury that can be missed due to inadequate initial and postoperative x-ray images and incorrect interpretation. ([10.1016/j.jse.2008.10.020](https://doi.org/10.1016/j.jse.2008.10.020))
- [L1] Timing of surgery did not affect Oxford Shoulder Score at any stage of follow-up, irrespective of age or fracture type. ([10.1302/0301-620x.102b1.bjj-2020-0546.r1](https://doi.org/10.1302/0301-620x.102b1.bjj-2020-0546.r1))

References

[1] Shoulder hemiarthroplasty for acute fractures of the proximal humerus: A minimum five-year follow-up. *Journal of Shoulder and Elbow Surgery*. 2008. DOI: [10.1016/j.jse.2007.06.025](https://doi.org/10.1016/j.jse.2007.06.025) [2] How the greater tuberosity affects clinical outcomes after reverse shoulder arthroplasty for proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2018. DOI: [10.1016/j.jse.2018.05.030](https://doi.org/10.1016/j.jse.2018.05.030) [3] Technique and clinical results of a new intramedullary support nail and plate system for fixation of 3- or 4- part proximal humeral fractures in older adults. *BMC Musculoskeletal Disorders*. 2022. DOI: [10.1186/s12891-022-05998-z](https://doi.org/10.1186/s12891-022-05998-z) [4] Mortality after inpatient

stay for proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2020. DOI: 10.1016/j.jse.2019.05.030 [5] Minimum 2-year outcomes of reverse total shoulder arthroplasty for fracture: how does acute arthroplasty compare with salvage?. *Journal of Shoulder and Elbow Surgery*. 2022. DOI: 10.1016/j.jse.2021.06.020 [6] Functional outcomes of reverse shoulder arthroplasty compared with hemiarthroplasty for acute proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2013. DOI: 10.1016/j.jse.2012.03.006 [7] Fracture line morphology of complex proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2017. DOI: 10.1016/j.jse.2017.05.014 [8] Improvement in shoulder rotation in complex shoulder fractures treated by reverse shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery*. 2013. DOI: 10.1016/j.jse.2012.03.011 [9] Ten-year follow-up of stemmed hemiarthroplasty for acute proximal humeral fractures. *The Bone & Joint Journal*. 2021. DOI: 10.1302/0301-620x.103b6.bjj-2020-1753.r1 [10] Readmissions, revisions, and mortality after treatment for proximal humeral fractures in three large states. *BMC Musculoskeletal Disorders*. 2019. DOI: 10.1186/s12891-019-2812-9 [11] Consensus statement on the treatment of proximal humerus fractures: a Delphi approach by the Neer Circle of the American Shoulder and Elbow Surgeons. *Journal of Shoulder and Elbow Surgery*. 2025. DOI: 10.1016/j.jse.2024.12.005 [12] Complications after non-surgical management of proximal humeral fractures: a systematic review of terms and definitions. *BMC Musculoskeletal Disorders*. 2019. DOI: 10.1186/s12891-019-2459-6 [13] Functional Outcome After Nonoperative Treatment of a Proximal Humeral Fracture in Adults. *Journal of Bone and Joint Surgery*. 2021. DOI: 10.2106/jbjs.20.02018 [14] Comparison of Hemiarthroplasty and Reverse Arthroplasty for Treatment of Proximal Humeral Fractures. *Journal of Bone and Joint Surgery*. 2013. DOI: 10.2106/jbjs.l.01115 [15] Morbidity and mortality of fragility proximal humerus fractures: a retrospective cohort study of patients presenting to a level one trauma center. *Journal of Shoulder and Elbow Surgery*. 2022. DOI: 10.1016/j.jse.2022.03.006 [16] Fragility of randomized controlled trials on treatment of proximal humeral fracture. *Journal of Shoulder and Elbow Surgery*. 2022. DOI: 10.1016/j.jse.2022.01.141 [17] Isolated avulsion fracture of the lesser tuberosity of the humerus in adolescents. A report of two cases. *The Journal of Bone & Joint Surgery*. 1995. DOI: 10.2106/00004623-199509000-00020 [19] Hospital Readmissions After Surgical Treatment of Proximal Humerus Fractures: Is Arthroplasty Safer Than Open Reduction Internal Fixation?. *Clinical Orthopaedics & Related Research*. 2014. DOI: 10.1007/s11999-014-3613-y [20] One Versus 3-Week Immobilization Period for Nonoperatively Treated Proximal Humeral Fractures. *Journal of Bone and Joint Surgery*. 2021. DOI: 10.2106/jbjs.20.02137 [21] In-hospital Complications Are More Likely to Occur After Reverse Shoulder Arthroplasty Than After Locked Plating for Proximal Humeral Fractures. *Clinical Orthopaedics & Related Research*. 2021. DOI: 10.1097/corr.0000000000001776 [22] A new clinical sign to detect radiologically occult greater tuberosity fractures. *Journal of Shoulder and Elbow Surgery*. 2024. DOI: 10.1016/j.jse.2023.07.044 [23] Proximal Humeral Fracture Treatment in Adults. *Journal of Bone and Joint Surgery*. 2014. DOI: 10.2106/jbjs.l.01293 [24] Proximal Humeral Fractures: “Damned If You Operate, and Damned If You Don’t”. *Journal of Bone and Joint Surgery*. 2019. DOI: 10.2106/jbjs.19.01109 [25] Mortality after a proximal humeral fracture. *The Bone & Joint Journal*. 2020. DOI: 10.1302/0301-620x.102b11.bjj-2020-0627.r1 [26] Hemiarthroplasty versus reverse shoulder arthroplasty for treatment of proximal humeral fractures: a meta-analysis. *Journal of Shoulder and Elbow Surgery*. 2016. DOI: 10.1016/j.jse.2015.08.030 [27] Short-term Complications for Proximal Humerus Fracture Surgery Have Decreased: An Analysis of the National Surgical Quality Improvement Program Database. *Clinical Orthopaedics & Related Research*. 2022. DOI: 10.1097/corr.0000000000002391 [28] The role of greater tuberosity healing in reverse shoulder arthroplasty: a finite element analysis. *Journal of Shoulder and Elbow Surgery*. 2020. DOI: 10.1016/j.jse.2019.07.022 [29] Does objective shoulder impairment explain patient-reported functional

outcome? A study of proximal humerus fractures. *Journal of Shoulder and Elbow Surgery*. 2011. DOI: 10.1016/j.jse.2010.06.005 [30] Neer Award 2006: Biomechanical assessment of inferior tuberosity placement during hemiarthroplasty for four-part proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2008. DOI: 10.1016/j.jse.2007.06.017 [35] Evaluation of the Constant score: which is the method to assess the objective strength?. *BMC Musculoskeletal Disorders*. 2019. DOI: 10.1186/s12891-019-2795-6 [37] Does fracture of the dominant shoulder have any effect on functional and quality of life outcome compared with the nondominant shoulder?. *Journal of Shoulder and Elbow Surgery*. 2015. DOI: 10.1016/j.jse.2014.10.006 [38] The assessment of scapular radiographs. *The Bone & Joint Journal*. 2013. DOI: 10.1302/0301-620x.95b8.30631 [40] Altered Glenohumeral Biomechanics in Proximal Humeral Fracture Malunion. *Journal of the American Academy of Orthopaedic Surgeons*. 2020. DOI: 10.5435/jaaos-d-20-00555 [43] Effects of shoulder position on axillary nerve positions during the split lateral deltoid approach. *Journal of Shoulder and Elbow Surgery*. 2009. DOI: 10.1016/j.jse.2008.12.001 [44] A morphovolumetric study of head malposition in proximal humeral fractures based on 3-dimensional computed tomography scans: the control volume theory. *Journal of Shoulder and Elbow Surgery*. 2018. DOI: 10.1016/j.jse.2017.12.004 [46] Cost-Minimization Analysis and Treatment Trends of Surgical and Nonsurgical Treatment of Proximal Humerus Fractures. *The Journal of Hand Surgery*. 2020. DOI: 10.1016/j.jhsa.2020.03.022 [47] Reverse shoulder arthroplasty or nothing for patients with displaced proximal humeral fractures: a randomized controlled trial. *Journal of Shoulder and Elbow Surgery*. 2024. DOI: 10.1016/j.jse.2024.02.023 [48] Operative versus non-operative treatment for 2-part proximal humerus fracture: A multicenter randomized controlled trial. *PLOS Medicine*. 2019. DOI: 10.1371/journal.pmed.1002855 [49] Classification of proximal humeral fractures based on a pathomorphologic analysis. *Journal of Shoulder and Elbow Surgery*. 2016. DOI: 10.1016/j.jse.2015.08.006 [50] Evaluation and Management of Pediatric Proximal Humerus Fractures. *Journal of the American Academy of Orthopaedic Surgeons*. 2015. DOI: 10.5435/jaaos-d-14-00033 [51] Contemporary Management of Proximal Humeral Fractures. *Journal of the American Academy of Orthopaedic Surgeons*. 2025. DOI: 10.5435/jaaos-d-24-01073 [52] Locking plate versus nonsurgical treatment for proximal humeral fractures: better midterm outcome with nonsurgical treatment. *Journal of Shoulder and Elbow Surgery*. 2011. DOI: 10.1016/j.jse.2011.01.025 [53] Osteoporosis does not affect bone mineral density change in the proximal humerus or the functional outcome after open reduction and internal fixation of unilateral displaced 3- or 4-part fractures at 12-month follow-up. *Journal of Shoulder and Elbow Surgery*. 2023. DOI: 10.1016/j.jse.2022.07.008 [54] Reverse Shoulder Arthroplasty to Treat Proximal Humerus Fracture Sequelae: A Review. *Journal of the American Academy of Orthopaedic Surgeons*. 2024. DOI: 10.5435/jaaos-d-23-00740 [55] Path analysis of factors for functional outcome at one year in 463 proximal humeral fractures. *Journal of Shoulder and Elbow Surgery*. 2011. DOI: 10.1016/j.jse.2011.06.009 [56] A comprehensive classification of proximal humeral fractures: HGLS system. *Journal of Shoulder and Elbow Surgery*. 2013. DOI: 10.1016/j.jse.2012.09.018 [57] CORR Insights®: What Factors Are Associated With Poor Shoulder Function and Serious Complications After Internal Fixation of Three-part and Four-part Proximal Humerus Fracture-dislocations?. *Clinical Orthopaedics & Related Research*. 2022. DOI: 10.1097/corr.0000000000002242 [58] Four-segment classification of proximal humeral fractures revisited: A multicenter study on 509 cases. *Journal of Shoulder and Elbow Surgery*. 2009. DOI: 10.1016/j.jse.2009.01.018 [59] Outcome measures in the management of proximal humeral fractures: a systematic review of their use and psychometric properties. *Journal of Shoulder and Elbow Surgery*. 2011. DOI: 10.1016/j.jse.2010.10.028 [60] Evaluation of the Neer System of Classification of Proximal Humeral Fractures with Computerized Tomographic Scans and Plain Radiographs. *The Journal of Bone & Joint Surgery*. 1996. DOI: 10.2106/00004623-199609000-00012 [62] *Outcomes after percutaneous reduction and fixation of proximal humeral fractures*. *Journal of Shoulder and Elbow*

Surgery. 2007. DOI: 10.1016/j.jse.2006.09.006 [63] Long-term outcomes of reverse shoulder arthroplasty versus nonoperative treatment for 3- or 4-part proximal humerus fractures in elderly patients: results from a prior randomized clinical trial. *Journal of Shoulder and Elbow Surgery*. 2025. DOI: 10.1016/j.jse.2024.09.032 [65] Is there still a place for anatomic hemiarthroplasty in patients with high functional demands in primary, nonreconstructable proximal humeral fractures? A clinical and radiographic assessment. *Journal of Shoulder and Elbow Surgery*. 2023. DOI: 10.1016/j.jse.2023.02.118 [66] Clinical outcomes of reverse total shoulder arthroplasty for elective indications versus acute 3- and 4-part proximal humeral fractures: a systematic review and meta-analysis. *Journal of Shoulder and Elbow Surgery*. 2022. DOI: 10.1016/j.jse.2021.07.014 [67] *Hemiarthroplasty for Humeral Four-part Fractures for Patients 65 Years and Older: A Randomized Controlled Trial*. *Clinical Orthopaedics & Related Research*. 2012. DOI: 10.1007/s11999-012-2531-0 [71] The effect of medial calcar support on proximal humeral fractures treated with locking plates. *Journal of Orthopaedic Surgery and Research*. 2022. DOI: 10.1186/s13018-022-03337-5 [72] Missed fractures of the greater tuberosity. *BMC Musculoskeletal Disorders*. 2018. DOI: 10.1186/s12891-018-2225-1 [73] Coronal displacement in proximal humeral fractures: correlation between shoulder radiographic and computed tomography scan measurements. *Journal of Shoulder and Elbow Surgery*. 2017. DOI: 10.1016/j.jse.2016.05.016 [74] CORR Insights®: 3D-printed Handheld Models Do Not Improve Recognition of Specific Characteristics and Patterns of Three-part and Four-part Proximal Humerus Fractures. *Clinical Orthopaedics & Related Research*. 2021. DOI: 10.1097/corr.0000000000002017 [76] Detection, classification, and characterization of proximal humerus fractures on plain radiographs. *The Bone & Joint Journal*. 2024. DOI: 10.1302/0301-620x.106b11.bjj-2024-0264.r1 [79] Missed posterior dislocation of the shoulder after intramedullary fixation of humeral fractures: A report of three cases. *Journal of Shoulder and Elbow Surgery*. 2009. DOI: 10.1016/j.jse.2008.10.020 [80] Does time to surgery affect patient-reported outcome in proximal humeral fractures? A subanalysis of the PROFHER randomized clinical trial. *The Bone & Joint Journal**. 2020. DOI: 10.1302/0301-620x.102b1.bjj-2020-0546.r1